

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

1

Find

(a)  $\int 3e^{4x+2} dx$

(b)  $\int (4e^{4-x} + 2) dx$

(c)  $\int \frac{e^{2x}+1}{4e^{-x}}$

TAP FOR ANSWERS

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

2

The functions  $f$  and  $g$  are defined by

$$f: x \rightarrow 5x + 2, x \in \mathbb{R}$$

$$g: x \rightarrow \frac{1}{x}, x \in \mathbb{R}, x \neq 0$$

(a) Find the following functions, stating the domain in each case.

(i)  $f^{-1}(x)$

(ii)  $fg(x)$

(iii)  $(fg)^{-1}(x)$

(b) Solve the equation  $f^{-1}(x) = fg(x)$ , giving your answers to 2 decimal places.

TAP FOR ANSWERS

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

3

A parachutist drops from a helicopter  $H$  and falls vertically from rest towards the ground. Her parachute opens 2 s after she leaves  $H$  and her speed then reduces to  $4 \text{ m s}^{-1}$ . For the first 2 s her motion is modelled as that of a particle falling freely under gravity. For the next 5 s the model is motion with constant deceleration, so that her speed is  $4 \text{ m s}^{-1}$  at the end of this period. For the rest of the time before she reaches the ground, the model is motion with constant speed of  $4 \text{ m s}^{-1}$ .

- (a) Sketch a speed-time graph to illustrate her motion from  $H$  to the ground.
- (b) Find her speed when the parachute opens.

A safety rule states that the helicopter must be high enough to allow the parachute to open and for the speed of a parachutist to reduce to  $4 \text{ m s}^{-1}$  before reaching the ground. Using the assumptions made in the above model,

- (c) find the minimum height of  $H$  for which the woman can make a drop without breaking this safety rule.

Given that  $H$  is 125 m above the ground when the woman starts her drop,

- (d) find the total time taken for her to reach the ground.
- (e) State one way in which the model could be refined to make it more realistic.

TAP FOR ANSWERS

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

4

(a) A particle rests in limiting equilibrium on a plane inclined at  $30^\circ$  to the horizontal.

Determine the acceleration with which the particle will slide down the plane when the angle of inclination is increased to  $45^\circ$ .

(b) A lift is accelerating upwards at  $1.5 \text{ m s}^{-2}$ . A girl of mass  $30 \text{ kg}$  is standing in the lift. Modeling the girl as a particle, find the force between her and the floor of the lift.

TAP FOR ANSWERS


# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

5

*Draw a force diagram and resolve forces for each block in appropriate directions.*

A block of 5kg rests on a slope which is  $30^\circ$  to the horizontal it is connected by a light inextensible string which passes through a frictionless pulley to a second block of 1kg. Find the tension in the string, the acceleration of the blocks and the direction of travel.



TAP FOR ANSWERS

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

6

An object moves in a straight line from a point  $O$ . at time  $t$  seconds the object has acceleration,  $a$ , where

$$a = -\cos 4\pi t \text{ m s}^{-2}, 0 \leq t \leq 4$$

When  $t = 0$ , the velocity of the object is  $0 \text{ m s}^{-1}$  and its displacement is  $0 \text{ m}$ .  
Find:

- (a) An expression for the velocity at time  $t$  seconds.
- (b) The maximum speed of the object
- (c) An expression for the displacement of the object at time  $t$  seconds.
- (d) The maximum displacement of the object from  $O$
- (e) The number of times the object changes direction during its motion.

TAP FOR ANSWERS

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

7

Two cyclists,  $C$  and  $D$ , are travelling with constant velocities  $(5\mathbf{i} - 2\mathbf{j}) \text{ m s}^{-1}$  and  $8\mathbf{j} \text{ m s}^{-1}$  respectively relative to a fixed origin  $O$ .

(a) Find the velocity of  $C$  relative to  $D$ .

At noon, the position vectors of  $C$  and  $D$  are  $(100\mathbf{i} + 300\mathbf{j}) \text{ m}$  and  $(150\mathbf{i} + 100\mathbf{j}) \text{ m}$  respectively, referred to  $O$ . At  $t$  seconds after noon, the position vector of  $C$  relative to  $D$  is  $\mathbf{s}$  metres.

(b) Show that  $\mathbf{s} = (-50 + 5t)\mathbf{i} + (200 - 10t)\mathbf{j}$ .

(c) By considering  $|\mathbf{s}|^2$ , or otherwise, find the value of  $t$  for which  $C$  and  $D$  are closest together.

TAP FOR ANSWERS

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

8

$P$  is the point  $(5, 6, -2)$ ,  $Q$  is the point  $(2, -1, 1)$  and  $R$  is the point  $(2, -3, 6)$ .

(a) Find the vectors  $\overrightarrow{PQ}$ ,  $\overrightarrow{PR}$  and  $\overrightarrow{QR}$

(b) Hence, or otherwise, find the area of triangle  $PQR$ .

TAP FOR ANSWERS



**BHASVIC MαTHS**  
**A1 DOUBLES ASSIGNMENT 19A**

9

Prove that the derivative of  $\cos x$  is  $-\sin x$  from first principles

TAP FOR ANSWERS

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

10

Liquid dye is poured onto a large flat cloth and forms a circular stain, the area of which grows at a steady rate of  $1.5 \text{ cm}^2 \text{ s}^{-1}$

Calculate, correct to 3 s.f.,

- (a) the radius, in cm, of the stain 4 seconds after it started forming
- (b) the rate, in  $\text{cm s}^{-1}$ , of increase of the radius of the stain after 4 seconds

TAP FOR ANSWERS

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

11

A curve has equation  $2x^2 + xy + y^2 = 14$

(a) Show clearly that  $\frac{dy}{dx} = -\frac{4x+y}{x+2y}$

(b) hence find the co-ordinates of the turning points of the curve.

TAP FOR ANSWERS

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

12

Evaluate the following


$$\lim_{\delta x \rightarrow 0} \sum_{x=0}^2 \frac{1}{\sqrt{4x+1}} dx$$

TAP FOR ANSWERS

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

13



The Diagram shows the graph of  $y = (1 + \sin 2x)^2$ ,  $0 \leq x \leq \frac{3\pi}{4}$

(a) Show that  $(1 + 2 \sin 2x)^2 \equiv \frac{1}{2}(3 + 4 \sin 2x - \cos 4x)$ .

(b) Hence find the area of the shaded region R.

(c) Find the coordinates of A, the turning point on the graph.

TAP FOR ANSWERS

# BHASVIC MαTHS


## A1 DOUBLES ASSIGNMENT 19A

14

The function  $f$  is defined by

$$f(x) = -\frac{5}{3}|x + 4| + 8, x \in \mathbb{R}$$

The diagram shows a sketch of the graph  $y = f(x)$



- State the range of  $f$ .
- Give a reason why  $f^{-1}(x)$  does not exist.
- Solve the inequality  $f(x) > \frac{2}{3}x + 4$ .
- State the range of values of  $k$  for which the equation  $f(x) = \frac{5}{3}x + k$  has no solutions

TAP FOR ANSWERS

# BHASVIC MATHS

## A1 DOUBLES ASSIGNMENT 19A

15

$$f(x) = x^3 - 2x - 1$$

- (a) Show that the equation  $f(x) = 0$  has a root,  $\alpha$ , at the interval  $1 < \alpha < 2$ .
- (b) Using  $x_0 = 1.5$  as a first approximation to  $\alpha$ , apply the Newton-Raphson procedure once to  $f(x)$  to find a second approximation to  $\alpha$ , giving your answer to 3 decimal places.

$$f(x) = x^2 - \frac{4}{x} + 6x - 10, x \neq 0$$

- c) Use differentiation to find  $f'(x)$ .

The root,  $\alpha$ , of the equation  $f(x) = 0$  lies in the interval  $[-0.4, -0.3]$

- (d) Taking  $-0.4$  as a first approximation to  $\alpha$ , apply the Newton-Raphson process once to  $f(x)$  to obtain a second approximation to  $\alpha$ . Give your answer to 3 decimal places.

TAP FOR ANSWERS

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

16

Complete this old spec paper

[https://www.madasmaths.com/archive/iygb\\_practice\\_papers/c3\\_practice\\_papers/c3\\_p.pdf](https://www.madasmaths.com/archive/iygb_practice_papers/c3_practice_papers/c3_p.pdf)

TAP FOR ANSWERS



# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

### 1 - Answers

(a)  $\frac{3}{4}e^{4x+2} + c$

(b)  $-4e^{4-x} + 2x + c$

(c)  $\frac{1}{12}e^{3x} + \frac{1}{4}e^x + c$

TAP TO RETURN

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

### 2 – Answers

(a) (i)  $f^{-1}: x \rightarrow \frac{x-2}{5}, x \in \mathbb{R}$

(ii)  $fg: x \rightarrow \frac{5}{x} + 2, x \in \mathbb{R}, x \neq 0$

(iii)  $(fg)^{-1}: x \rightarrow \frac{5}{x-2}, x \in \mathbb{R}, x \neq 2$

(b) 13.81, -1.81

TAP TO RETURN

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

### 3 - Answers

(b)  $19.6\text{ms}^{-1}$

(c) 78.6m

(d) 18.6 seconds

TAP TO RETURN

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

### 4 - Answers

(a)  $2.0 \text{ ms}^{-2}$  (2sf)

(b) 340 N (2sf)

TAP TO RETURN

# BHAVIC MATHS

## A1 DOUBLES ASSIGNMENT 19A

### 5 - Answers

$T=12.25\text{N}$  (3sf),  $a= 2.45 \text{ ms}^{-2}$  down the plane for the 5kg block,

TAP TO RETURN

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

### 6 - Answers

(a)  $v = -\frac{\sin 4\pi t}{4\pi}$

(b)  $\frac{1}{4\pi}$

(c)  $s = \frac{\cos 4\pi t}{16\pi^2} - \frac{1}{16\pi^2}$

(d)  $\frac{1}{8\pi^2}$

(e) 16

TAP TO RETURN

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

### 7 - Answers

(a)  $5\mathbf{i} - 10\mathbf{j} \text{ m s}^{-1}$

(b)  $\sqrt{10}$

(c)  $5\mathbf{i} - 10\mathbf{j} \text{ m s}^{-1}$

(d)  $t = 18$

TAP TO RETURN

**BHASVIC MαTHS**  
**A1 DOUBLES ASSIGNMENT 19A**

8 - Answers

(a)  $\overrightarrow{PQ} = -3\mathbf{i} - 7\mathbf{j} + 3\mathbf{k}$ ,  $\overrightarrow{PR} = -3\mathbf{i} - 9\mathbf{j} + 8\mathbf{k}$ ,  $\overrightarrow{QR} = -2\mathbf{j} + 5\mathbf{k}$

(b) 17

TAP TO RETURN



**BHASVIC MαTHS**  
**A1 DOUBLES ASSIGNMENT 19A**

9 - Answers

Proof

TAP TO RETURN

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

### 10 - Answers

(a) 1.38 cm

(b)  $0.173 \text{ cm s}^{-1}$

TAP TO RETURN

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

### 11 - Answers

(1,-4) and (-1,4)

TAP TO RETURN

**BHASVIC MαTHS**  
**A1 DOUBLES ASSIGNMENT 19A**

12 - Answers

1

TAP TO RETURN

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

### 13 - Answers

$$\begin{aligned} \text{(a)} \quad (1 + \sin 2x)^2 &\equiv 1 + 2 \sin 2x + \sin^2 2x \\ &\equiv 1 + 2 \sin 2x + \frac{1 - \cos 4x}{2} \equiv \frac{3}{2} + 2 \sin 2x - \frac{\cos 4x}{2} \\ &\equiv \frac{1}{2} (3 + 4 \sin 2x - \cos 4x) \end{aligned}$$

$$\text{(b)} \quad \frac{9\pi}{8} + 1$$

$$\text{(c)} \quad \left(\frac{\pi}{4}, 4\right)$$

TAP TO RETURN

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

### 14 - Answers

(a)  $f(x) \leq 8$

(b) The function is not one-to-one

(c)  $-\frac{32}{3} < x < -\frac{8}{7}$

(d)  $k > \frac{44}{3}$

TAP TO RETURN

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

### 15 - Answers

(a)  $f(1) = -2, f(2) = 3$  There is a sign change in the interval  $1 < \alpha < 2$ , so there is a root in this interval.

(b)  $x_1 = 1.632$

(c)  $f'(x) = 2x + \frac{4}{x^2} + 6$

(d) 0.326

TAP TO RETURN

# BHASVIC MαTHS

## A1 DOUBLES ASSIGNMENT 19A

### 16 - Answers

- (a) [https://www.madasmaths.com/archive/iygb\\_practice\\_papers/c3\\_practice\\_papers/c3\\_p\\_solutions.pdf](https://www.madasmaths.com/archive/iygb_practice_papers/c3_practice_papers/c3_p_solutions.pdf)

TAP TO RETURN