

Question		Done	BP	Ready	Topic	Comment
Drill	Aa				C4 Implicit Diff – $xy = \sin^2$	$\frac{y}{\sin 2y - x}$
	Ab				C4 Implicit Diff – $\tan(2x + y) = x$	$\cos^2(2x + y) - 2$
	Ac				C4 Implicit Diff – $e^{(xy)} = 4$	$-\frac{y}{x}$
	Ba				C3 Functions – MOD sketch	Check using autograph/desmos
	Bb				C3 Functions – MOD sketch	Check using autograph/desmos
	Bc				C3 Functions – MOD sketch	Check using autograph/desmos
	Ca				C4 Parametric – dy/dx chain rule	$\frac{1}{t}$
	Cb				C4 Parametric – dy/dx chain rule	$-\frac{2}{5}t^2$
	Cc				C4 Parametric – dy/dx chain rule	$\cot t$
	Da				C4 Vectors – Find cosine acute angle	$\frac{13}{\sqrt{14}\sqrt{26}}$
	Db				C4 Vectors – Find cosine acute angle	$\frac{3}{7\sqrt{2}}$
	Dc				C4 Vectors – Find cosine acute angle	$\frac{1}{6}$ *remember it's the acute angle! So mod signs! *
Applied consolidation	1a				M2 Projectiles – cliff 50m time of flight	5 s
	1b				M2 Projectiles – horizontal landing dist	130m
	1c				M2 Projectiles – speed & direction impact	43ms^{-1} , 53° to the horizontal
	1d				M2 Projectiles – show after T travels 45 deg	Proof
	2a				M2 Kinematics – given F, vectors, find acc	$14\mathbf{i} + 18\mathbf{j} \text{ ms}^{-2}$
	2b				M2 Kinematics – given F, vectors, find sp	37.4 ms^{-1}
	3				M2 Power – find max acc given power	450N , $\frac{9}{16} \text{ ms}^{-2}$
	4a				M2 Power – find power given up slope	50.4kW
	4b				M2 Power – find max speed up slope	95 kmh^{-1}
	5a				C3 Trig – Rmethod, involving 2x	e.g. if using Rcos $R = 13$, $\alpha = 1.176$
Core consolidation	5b				C3 Trig – solve using R method	1.13, 0.0425
	5ci)				C3 Trig – R method write f(x)max	$f(x) = 13$
	5cii)				C3 Trig – R method smallest +ve value x	$x = 0.588$
	6a				C3 Functions – show f(x) rearrange	show
	6b				C3 Functions – find range f	$0 < f(x) < 1/4$
	6c				C3 Functions – find f^{-1} and state its domain	$f^{-1}(x) = \frac{1}{x} - 1$, $0 < x < \frac{1}{4}$
	6d				C3 Functions – solve $fg(x) = 1/8$	$x = \pm\sqrt{5}$
	7ai)				C3 Diff – product rule	$e^{3x}(\sin x + 7 \cos x)$

7aii)			C3 Diff – product rule involving ln	$3x^2 \ln(5x + 2) + \frac{5x^3}{5x + 2}$
7b			C3 Diff – quotient rule	
7c			C3 Diff – second derivative & solve	$x = 1, -3$
8a			C4 Integral – Trapezium rule complete table	$e^{0.32}, e^{1.28}$
8b			C4 Integral – use trapezium rule	4.922
9			C4 Connected rates – balloon inflating	$\frac{5k}{3}$
10a			C4 Diff Eq – partial fractions	$\frac{\frac{1}{2}}{(2 - y)} + \frac{\frac{1}{2}}{(2 + y)}$
10b			C4 Diff Eq – find particular solution	$\sec^2 x = \frac{8 + 4y}{2 - y}$
11			C4 Vectors – shortest dist from point to L	$2\sqrt{46}$ * should get $\lambda = 0$ when dotting line with AX line segment *
12			C4 Connected Rates – Area between circles	$\frac{dA}{dt} = -8\pi \text{ cm}^2\text{s}^{-1}$

α	β	γ	δ	ε	ζ	η	θ	ι	κ	λ	μ	ν	ξ	\omicron	π	ρ	σ	τ	υ	ϕ	χ	ψ	ω
----------	---------	----------	----------	---------------	---------	--------	----------	---------	----------	-----------	-------	-------	-------	------------	-------	--------	----------	--------	------------	--------	--------	--------	----------

"It is not knowledge but the act of learning which grants the greatest enjoyment" K. F. Gauss

A2 Maths with Mechanics Assignment τ (tau)

Due in week beginning 26th February

Drill

Part A Find $\frac{dy}{dx}$ for each of the following functions:

(a) $xy = \sin^2 y$ (b) $\tan(2x + y) = x$ (c) $e^{xy} = 4$

Part B Sketch the following functions:

(a) $y = 1 - |x + 2|$ (b) $y = 3|\sin x|$ (c) $y = 2\ln|x| + 2$

Part C Find $\frac{dy}{dx}$ using the chain rule, for each of the following functions:

(a) $x = t^2 + 1$
 $y = 2t$ (b) $x = \frac{5}{t}$
 $y = 2t$ (c) $x = 1 - \cos t$
 $y = \sin t$

Part D Find the cosine of the acute angle between the following pairs of vectors:

(a) $\mathbf{a} = 2\mathbf{i} - \mathbf{j} + 3\mathbf{k}$, $\mathbf{b} = 5\mathbf{i} + \mathbf{k}$
(b) $\mathbf{a} = 6\mathbf{i} - 3\mathbf{j} + 2\mathbf{k}$, $\mathbf{b} = \mathbf{i} + \mathbf{j}$
(c) $\mathbf{a} = \mathbf{i} + \mathbf{j} - 2\mathbf{k}$, $\mathbf{b} = 2\mathbf{i} - \mathbf{j} + \mathbf{k}$

Mechanics Consolidation

- A stone is thrown up at an angle of 30° to the horizontal with a speed of 30 ms^{-1} from the edge of a cliff 50 m above sea level. If the stone lands in the sea, calculate:
 - How long it is in the air,
 - How far from the base of the cliff it lands,
 - The speed and direction of the stone as it hits the water.
 - Show that after T secs it will be travelling at an angle 45° below the horizontal.

$$\text{where } T = \frac{15(\sqrt{3} + 1)}{g}$$

2. The resultant force \mathbf{F} Newtons acting on a particle of mass 0.5kg at time t s is given by: $\mathbf{F} = (t^2 - 2)\mathbf{i} + (2t + 3)\mathbf{j}$
- a) Find an expression for the acceleration of the particle after 3 secs.
- b) Given that the velocity of the particle at time $t = 0$ is $4\mathbf{i} \text{ ms}^{-1}$, find the speed after 3 secs.
3. A car has an engine of maximum power 15kW. Calculate the force resisting the motion of the car when it is travelling at its maximum speed of 120 kmh^{-1} on a level road. Assuming an unchanged resistance and taking the mass of the car to be 800kg, calculate the maximum acceleration of the car when travelling at 60 kmh^{-1} on a level road.
(Note: UNITS!)
4. A car of mass 1 tonne is moving at a constant velocity of 60 km per hour up an inclined road which makes an angle of 6° with the horizontal.
- a) Given that the non-gravitational resistance down the slope is 2000N, find the rate at which the car is working,
- b) If the engine has a maximum power output of 80kw, calculate the maximum speed of the car up the same slope.

Core consolidation

5. $f(x) = 5 \cos 2x + 12 \sin 2x$.
- (a) Using a suitable R-Method involving Rcos or Rsin, find the value of R and the value of α to 3 decimal places.
- (b) Hence solve the equation
- $$5 \cos 2x + 12 \sin 2x = 6 \quad \text{for } 0 \leq x < \pi.$$
- (c) (i) Write down the maximum value of $f(x)$
- (ii) Find the smallest positive value of x for which this maximum value occurs.

6. The function f is defined by

$$f: x \mapsto \frac{2(x-1)}{x^2-2x-3} - \frac{1}{x-3}, \quad x > 3.$$

- (a) Show that $f(x) = \frac{1}{x+1}$, $x > 3$.
- (b) Find the range of f .
- (c) Find $f^{-1}(x)$. State the domain of this inverse function.

The function g is defined by

$$g: x \mapsto 2x^2 - 3, \quad x \in \mathbb{R}.$$

(d) Solve $fg(x) = \frac{1}{8}$.

7. (a) Differentiate with respect to x ,

(i) $e^{3x}(\sin x + 2 \cos x)$,

(ii) $x^3 \ln(5x + 2)$.

Given that $y = \frac{3x^2 + 6x - 7}{(x+1)^2}$, $x \neq -1$,

(b) Show that $\frac{dy}{dx} = \frac{20}{(x+1)^3}$.

(c) Hence find $\frac{d^2y}{dx^2}$ and the real values of x for which $\frac{d^2y}{dx^2} = -\frac{15}{4}$.

8.




Figure 1

Figure 1 shows part of the curve with equation $y = e^{0.5x^2}$. The finite region R , shown shaded in Figure 1, is bounded by the curve, the x -axis, the y -axis and the line $x = 2$.

- (a) Copy and complete the table with the values of y corresponding to $x = 0.8$ and $x = 1.6$.

x	0	0.4	0.8	1.2	1.6	2
y	e^0	$e^{0.08}$		$e^{0.72}$		e^2

- (b) Use the trapezium rule with all the values in the table to find an approximate value for the area of R , giving your answer to 4 significant figures.

9. A spherical balloon is being blown up at a rate proportional to its volume at the time. Given the volume of a sphere is $\frac{4}{3}\pi r^3$, find the rate of change of the radius of the balloon in terms of k at the moment when the radius is 5cm.

10. (a) Express $\frac{2}{4-y^2}$ in partial fractions.

- (b) Hence obtain the solution of

$$2 \cot x \frac{dy}{dx} = (4 - y^2)$$

for which $y = 0$ at $x = \frac{\pi}{3}$, giving your answer in the form $\sec^2 x = g(y)$.

11. There is a line with equation $\underline{r} = (4\underline{i} - 3\underline{j} - 7\underline{k}) + \lambda(3\underline{i} - 3\underline{j} + 2\underline{k})$. A has position vector $(2\underline{i} + 3\underline{j} + 5\underline{k})$, find the shortest distance from the line to A.

12 At a given instant, the radii of two concentric circles are 8cm and 12cm. The radius of the outer circle is increasing at a rate of 1cm s^{-1} , and the radius of the inner circle is increasing at a rate of 2cm s^{-1} . Find the rate of change of the area enclosed by the two circles at that instant.